arXiv:math/0512181v1 [math.ST] 8 Dec 2005

Accompanying document to "Point Estimation with Exponentially Tilted Empirical Likelihood"

Susanne M. Schennach

First draft: December 12, 2002
This version: December 7, 2005

This manuscript provide the calculational details involved in the proof of the equivalence between the $O\left(n^{-2}\right)$ variance of the ETEL and EL estimator that are omitted from the appendix of "Point Estimation with Exponentially Tilted Empirical Likelihood" by Susanne M. Schennach.
Whenever possible, the following calculations were verified using the symbolic calculations capabilities of Maple.

1 Definitions

Let

$$\begin{aligned}\beta &= \left(\tau,\kappa',\lambda',\theta'\right)' \\ \beta^* &= \left(1,\mathbf{0}',\mathbf{0}',\theta^{*\prime}\right)'\end{aligned}$$

$$\begin{aligned}\dot{g} &= g\left(x_i,\theta\right) \\ \dot{G} &= \frac{\partial g\left(x_i,\theta\right)}{\partial\theta'}\end{aligned}$$

To simplify the notation, the dependence of the above quantities on i and θ is implicit.

$$\begin{aligned}G &= E\left[G\left(x_i,\theta^*\right)\right] \\ \Omega &= E\left[g\left(x_i,\theta^*\right)g\left(x_i,\theta^*\right)'\right]\end{aligned}$$

$$\begin{aligned}\bar{g} &= \left[n^{-1/2}\sum_i \dot{g}\right]_{\beta=\beta^*} \\ \bar{G} &= \left[n^{-1/2}\sum_i\left(\dot{G}-G\right)\right]_{\beta=\beta^*} \\ \bar{\Omega} &= \overline{gg'} = \left[n^{-1/2}\sum_i\left(\dot{g}\dot{g}'-\Omega\right)\right]_{\beta=\beta^*}\end{aligned}$$

Convention for transpose: $G'_{\cdot h}\equiv\left(G_{\cdot h}\right)'$
Expectations evaluated at $\beta=\beta^*$ will be denoted by $E^*[\ldots]$. For instance, $E^*\left[\dot{G}\right]=G$ and $E^*\left[\dot{g}\dot{g}'\right]=\Omega$.

1.1 Moment conditions for ETEL

In ETEL, $\hat{\beta}^{ETEL}$ solves $n^{-1}\sum_i\dot{\phi}^{ETEL}=0$, where

$$\dot{\phi}^{ETEL} = \begin{bmatrix} \dot{\tau} - \tau \\ \dot{\tau}\dot{g} \\ \left(\tau - \dot{\tau}\right)\dot{g} + \dot{\tau}\dot{g}\dot{g}'\kappa \\ \dot{\tau}\dot{G}'\kappa + \dot{\tau}\dot{G}'\lambda\dot{g}'\kappa - \dot{\tau}\dot{G}'\lambda + \tau\dot{G}'\lambda \end{bmatrix}$$

and where

$$\dot{\tau} = \exp\left(\lambda'\dot{g}\right).$$

Define

$$\begin{aligned} \Phi^{ETEL}_{l,j} &= E^*\left[\frac{\partial\dot{\phi}^{ETEL}_l}{\partial\beta_j}\right] \\ \Phi^{ETEL}_{l,jk} &= E^*\left[\frac{\partial^2\dot{\phi}^{ETEL}_l}{\partial\beta_j\partial\beta_k}\right] \\ \Phi^{ETEL}_{l,jkh} &= E^*\left[\frac{\partial^3\dot{\phi}^{ETEL}_l}{\partial\beta_j\partial\beta_k\partial\beta_h}\right] \end{aligned}$$

$$\begin{aligned} \bar{\Phi}^{ETEL}_{l} &= \left[n^{-1/2}\sum_i \dot{\phi}^{ETEL}_l\right]_{\beta=\beta^*} \\ \bar{\Phi}^{ETEL}_{l,j} &= \left[n^{-1/2}\sum_i \left(\frac{\partial\dot{\phi}^{ETEL}_l}{\partial\beta_j} - \Phi^{ETEL}_{l,j}\right)\right]_{\beta=\beta^*} \\ \bar{\Phi}^{ETEL}_{l,jk} &= \left[n^{-1/2}\sum_i \left(\frac{\partial^2\dot{\phi}^{ETEL}_l}{\partial\beta_j\partial\beta_k} - \Phi^{ETEL}_{l,jk}\right)\right]_{\beta=\beta^*} \\ \bar{\Phi}^{ETEL}_{l,jkh} &= \left[n^{-1/2}\sum_i \left(\frac{\partial^3\dot{\phi}^{ETEL}_l}{\partial\beta_j\partial\beta_k\partial\beta_h} - \Phi^{ETEL}_{l,jkh}\right)\right]_{\beta=\beta^*} \end{aligned}$$

$$\begin{aligned} \bar{\Phi}^{ETEL}_{\cdot,\cdot} &= \begin{bmatrix} -1 & 0 & \bar{g}' & 0 \\ 0 & 0 & \bar{\Omega} & \bar{G} \\ \bar{g} & \bar{\Omega} & -\bar{\Omega} & 0 \\ 0 & \bar{G}' & 0 & 0 \end{bmatrix}_{\beta=\beta^*} \\ \Phi^{ETEL}_{\cdot,\cdot} &= \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & \Omega & G \\ 0 & \Omega & -\Omega & 0 \\ 0 & G' & 0 & 0 \end{bmatrix} \end{aligned}$$

where the dot replacing the subscripts denote all the elements of the matrix.

1.2 Moment conditions for EL

The moment conditions for the EL parameters $\left(\hat{\kappa}', \hat{\theta}'\right)'$ are

$$n^{-1}\sum_i \begin{bmatrix} \dot{\varepsilon}\dot{g} \\ \dot{\varepsilon}\dot{G}'\kappa \end{bmatrix} = 0$$

where $\dot{\varepsilon} = \left(1 - \kappa'\dot{g}\right)^{-1}$ and κ is the Lagrange multiplier of the moment contraints, which has been relabelled κ to simplify the comparison with ETEL. Furthermore, to again simplify the comparison with ETEL, we augment this vector by $1+\dim\kappa$ additional moment conditions and introduce the same number of additional parameters (τ,λ) where $\tau\in\mathbb{R}$ and $\lambda\in\mathbb{R}^{\dim\kappa}$:

$$n^{-1}\sum_i \begin{bmatrix} \dot{\tau}-\tau \\ \dot{\tau}\dot{g} \\ \dot{\varepsilon}\dot{g} \\ \dot{\varepsilon}\dot{G}'\kappa \end{bmatrix} = 0$$

where $\dot{\tau}=\exp(\lambda'\dot{g})$. The additional moment conditions merely define the values of the new parameters $\left(\hat{\tau},\hat{\lambda}\right)$ and do not change the values of $\left(\hat{\kappa}',\hat{\theta}'\right)$. Indeed, whenever $\left(\hat{\kappa}',\hat{\theta}'\right)'$ are such that the bottom two subvectors are zero, one can always find a value of $\left(\hat{\tau},\hat{\lambda}\right)$ that will make the top two subvectors vanish as well. Since the origin is in the convex hull of $\left\{g\left(x_i,\hat{\theta}\right)\right\}_{i=1}^{n}$ w.p.a. 1, there exists $\hat{\lambda}$ such that $n^{-1}\sum_i\dot{\tau}\dot{g}=0$ w.p.a. 1. Then, we can just set $\hat{\tau}=n^{-1}\sum_i\dot{\tau}$.

Finally, since just-identified GMM is invariant under linear transformations of the vector of moment conditions, the moment conditions for EL can equivalently be written as $n^{-1}\sum_i\dot{\phi}^{EL}=0$, where

$$\dot{\phi}^{EL} = \begin{bmatrix} \dot{\tau}-\tau \\ \dot{\tau}\dot{g} \\ \dot{\varepsilon}\dot{g}-\dot{\tau}\dot{g} \\ \dot{\varepsilon}\dot{G}'\kappa \end{bmatrix}$$

This particular version of the EL moment conditions will drastically simplify our calculations, due to the fact that the matrices of first derivatives for ETEL and EL become nearly identical:

$$\begin{aligned}
\bar{\Phi}^{EL}_{\cdot,\cdot} &= \begin{bmatrix} -1 & 0 & \bar{g}' & 0 \\ 0 & 0 & \bar{\Omega} & \bar{G} \\ 0 & \bar{\Omega} & -\bar{\Omega} & 0 \\ 0 & \bar{G}' & 0 & 0 \end{bmatrix} \\
\Phi^{EL}_{\cdot,\cdot} &= \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & \Omega & G \\ 0 & \Omega & -\Omega & 0 \\ 0 & G' & 0 & 0 \end{bmatrix}
\end{aligned}$$

Note that

$$\Phi^{ETEL}_{\cdot,\cdot} = \Phi^{EL}_{\cdot,\cdot} \equiv \Phi_{\cdot,\cdot}$$

while $\bar{\Phi}^{ETEL}_{\cdot,\cdot}$ and $\bar{\Phi}^{EL}_{\cdot,\cdot}$ differ by a single element.

1.3 Other definitions and conventions

$$\begin{aligned}
\Psi^{ETEL}_{\cdot,j} &= -\left(\Phi_{\cdot,\cdot}\right)^{-1}\Phi^{ETEL}_{\cdot,j} \\
\Psi^{ETEL}_{\cdot,jk} &= -\left(\Phi_{\cdot,\cdot}\right)^{-1}\Phi^{ETEL}_{\cdot,jk} \\
\Psi^{ETEL}_{\cdot,jkh} &= -\left(\Phi_{\cdot,\cdot}\right)^{-1}\Phi^{ETEL}_{\cdot,jkh}
\end{aligned}$$

$$\begin{aligned}
\bar{\Psi}^{ETEL}_{\cdot} &= -\left(\Phi_{\cdot,\cdot}\right)^{-1}\bar{\Phi}^{ETEL}_{\cdot} \\
\bar{\Psi}^{ETEL}_{\cdot,j} &= -\left(\Phi_{\cdot,\cdot}\right)^{-1}\bar{\Phi}^{ETEL}_{\cdot,j} \\
\bar{\Psi}^{ETEL}_{\cdot,jk} &= -\left(\Phi_{\cdot,\cdot}\right)^{-1}\bar{\Phi}^{ETEL}_{\cdot,jk}
\end{aligned}$$

and similarly for EL.

Again, a dot replacing a given subscript denotes a vector of all the values taken for all values of that subscript.

e.g. $\Psi^{ETEL}_{\cdot,j} = \begin{bmatrix} \Psi^{ETEL}_{1,j} \\ \vdots \\ \Psi^{ETEL}_{1+2\dim\lambda+\dim\theta,j} \end{bmatrix}$, $\Psi^{ETEL}_{l,\cdot} = \begin{bmatrix} \Psi^{ETEL}_{l,1} & \cdots & \Psi^{ETEL}_{l,1+2\dim\lambda+\dim\theta} \end{bmatrix}$.

Following Newey and Smith (2001), define

$$\begin{aligned} P &= \Omega^{-1} - \Omega^{-1} G \left(G'\Omega^{-1}G\right)^{-1} G'\Omega^{-1} \\ H &= \left(G'\Omega^{-1}G\right)^{-1} G'\Omega^{-1} \\ \Sigma &= \left(G'\Omega^{-1}G\right)^{-1} \end{aligned}$$

Identities:

$$\begin{aligned} PG &= 0 \\ P' &= P \\ P\Omega P &= P \\ P\Omega H' &= 0 \\ H\Omega H' &= \Sigma \end{aligned}$$

Define

$$\begin{aligned} l_\tau &= 0 \\ l_\kappa &= 1 \\ l_\lambda &= 1 + \dim\kappa \\ l_\theta &= 1 + 2\dim\kappa \end{aligned}$$

These symbols will be used to isolate subvectors and submatrices. For instance,

$$\Phi^{EL}_{h,l_\kappa+j} = E\left[\frac{\partial \dot{\phi}^{EL}_h}{\partial \kappa_j}\right]$$

with j implicitly ranging from 1 to $\dim\kappa$.

2 Stochastic expansion

The following merely rewrites the conclusion of Lemma A4 in Newey and Smith (2001).using our notation.

$$\left(\hat{\beta} - \beta^*\right) = n^{-1/2}\bar{\Psi}_{\cdot} + n^{-1}\bar{Q}_{\cdot} + n^{-3/2}\bar{R}_{\cdot} + O_p\left(n^{-2}\right)$$

where

$$\begin{aligned} \bar{Q}_l &= \bar{\Psi}_{l,j}\bar{\Psi}_j + \frac{1}{2}\Psi_{l,jk}\bar{\Psi}_j\bar{\Psi}_k \\ \bar{R}_l &= \bar{\Psi}_{l,j}\bar{Q}_j + \Psi_{l,jk}\bar{Q}_j\bar{\Psi}_k + \frac{1}{2}\bar{\Psi}_{l,jk}\bar{\Psi}_j\bar{\Psi}_k + \frac{1}{6}\Psi_{l,jkh}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_h \end{aligned}$$

using the convention that repeated indices are summed over, e.g. $\bar{\Psi}_{l,j}\bar{\Psi}_j \equiv \sum_j \bar{\Psi}_{l,j}\bar{\Psi}_j$.
(We also simplified Newey and Smith's result using the fact that $\Psi_{l,jk} = \Psi_{l,kj}$).
The quantities associated with each estimator will distinguished by an "ETEL" or "EL" superscript.
The variance of $\hat{\beta}$ is:

$$\operatorname{Var}\left[\left(\hat{\beta} - \beta^*\right)\right] = n^{-1}\operatorname{Var}\left[\bar{\Psi}_{\cdot}\right] + n^{-2}\operatorname{Var}\left[\bar{Q}_{\cdot}\right] + n^{-2}\text{Covar}\left[\bar{R}_{\cdot}, \bar{\Psi}'_{\cdot}\right] + n^{-2}\text{Covar}\left[\bar{\Psi}_{\cdot}, \bar{R}'_{\cdot}\right] + o\left(n^{-2}\right)$$

3 Partitioned Inverse of $\Phi_{\cdot,\cdot}$

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & \Omega & G \\ 0 & \Omega & -\Omega & 0 \\ 0 & G' & 0 & 0 \end{bmatrix}^{-1} = \begin{bmatrix} -1 & \begin{bmatrix} 0 & 0 & 0 \end{bmatrix} \\ \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix} & \begin{bmatrix} 0 & \Omega & G \\ \Omega & -\Omega & 0 \\ G' & 0 & 0 \end{bmatrix}^{-1} \end{bmatrix}$$

where $\begin{bmatrix} 0 & \Omega & G \\ \Omega & -\Omega & 0 \\ G' & 0 & 0 \end{bmatrix}^{-1}$ can be found using partitioned inverse formula:

$$\begin{bmatrix} A_{11} & A_{12} \\ A_{21} & A_{22} \end{bmatrix}^{-1} = \begin{bmatrix} B_{11} & B_{12} \\ B_{21} & B_{22} \end{bmatrix}$$

$$A_{11}^{-1} = \begin{bmatrix} 0 & \Omega \\ \Omega & -\Omega \end{bmatrix}^{-1} = \begin{bmatrix} \Omega^{-1} & \Omega^{-1} \\ \Omega^{-1} & 0 \end{bmatrix}$$

$$B_{22} = \left(A_{22} - A_{21}A_{11}^{-1}A_{12}\right)^{-1} = \left(0 - \begin{bmatrix} G' & 0 \end{bmatrix} \begin{bmatrix} \Omega^{-1} & \Omega^{-1} \\ \Omega^{-1} & 0 \end{bmatrix} \begin{bmatrix} G \\ 0 \end{bmatrix}\right)^{-1} = -\left(G'\Omega^{-1}G\right)^{-1} \equiv -\Sigma$$

$$B_{11} = A_{11}^{-1}\left(I + A_{12}B_{22}A_{21}A_{11}^{-1}\right) = \begin{bmatrix} \Omega^{-1} & \Omega^{-1} \\ \Omega^{-1} & 0 \end{bmatrix} \left(I - \begin{bmatrix} G \\ 0 \end{bmatrix} \Sigma \begin{bmatrix} G' & 0 \end{bmatrix} \begin{bmatrix} \Omega^{-1} & \Omega^{-1} \\ \Omega^{-1} & 0 \end{bmatrix}\right) =$$

$$= \begin{bmatrix} \Omega^{-1} & \Omega^{-1} \\ \Omega^{-1} & 0 \end{bmatrix} \begin{bmatrix} I - G\Sigma G'\Omega^{-1} & -G\Sigma G'\Omega^{-1} \\ 0 & I \end{bmatrix} = \begin{bmatrix} \Omega^{-1} - \Omega^{-1}G\Sigma G'\Omega^{-1} & \Omega^{-1} - \Omega^{-1}G\Sigma G'\Omega^{-1} \\ \Omega^{-1} - \Omega^{-1}G\Sigma G'\Omega^{-1} & -\Omega^{-1}G\Sigma G'\Omega^{-1} \end{bmatrix}$$

$$= \begin{bmatrix} P & P \\ P & P - \Omega^{-1} \end{bmatrix}$$

$$B_{12} = -A_{11}^{-1}A_{12}B_{22} = -\begin{bmatrix} \Omega^{-1} & \Omega^{-1} \\ \Omega^{-1} & 0 \end{bmatrix} \begin{bmatrix} G \\ 0 \end{bmatrix} \Sigma = -\begin{bmatrix} \Omega^{-1}G\Sigma \\ \Omega^{-1}G\Sigma \end{bmatrix} = \begin{bmatrix} H' \\ H' \end{bmatrix}$$

$$B_{21} = B_{12}'$$

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 0 & \Omega & G \\ 0 & \Omega & -\Omega & 0 \\ 0 & G' & 0 & 0 \end{bmatrix}^{-1} = \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & P & P & H' \\ 0 & P & P - \Omega^{-1} & H' \\ 0 & H & H & -\Sigma \end{bmatrix}$$

For notational convenience, we define

$$\Phi_{jk}^{-1} \equiv \left(\Phi_{\cdot,\cdot}^{-1}\right)_{jk}$$

4 $\bar{\Psi}_\cdot$ Term

$$\bar{\Psi}_\cdot^{ETEL} = \bar{\Psi}_\cdot^{EL} \equiv \bar{\Psi}_\cdot = -\Phi_{\cdot,\cdot}^{-1}\bar{\Phi}_\cdot = -\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & P & P & H' \\ 0 & P & P - \Omega^{-1} & H' \\ 0 & H & H & -\Sigma \end{bmatrix} \begin{bmatrix} 0 \\ \bar{g} \\ 0 \\ 0 \end{bmatrix} = \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix}$$

$$\text{Var}\left(\bar{\Psi}_\cdot\right) = E\left[\begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} \begin{bmatrix} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{bmatrix}\right] = \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & P & P & 0 \\ 0 & P & P & 0 \\ 0 & 0 & 0 & \Sigma \end{bmatrix}$$

5 $\bar{Q}_\cdot$ TERM

$$\bar{Q}_l = \bar{\Psi}_{l,j}\bar{\Psi}_j + \bar{\Psi}_j\Psi_{l,jk}\bar{\Psi}_k/2$$

5.1 Proof that $\bar{Q}_l^{ETEL} - \bar{Q}_l^{EL} = 0$

This result will imply that all the parameters $\left(\hat{\tau}, \hat{\kappa}', \hat{\lambda}', \hat{\theta}\right)$ for ETEL and EL differ by less than $O_p\left(n^{-1}\right)$. This may come as a surprise since the Lagrange multipliers of EL and ET are known to differ by $O_p\left(n^{-1}\right)$. However, we have carefully defined the parameter vector of each estimator so that the Lagrange multiplier of EL ($\hat{\kappa}$) corresponds to the auxiliary parameter $\hat{\kappa}$ of ETEL instead of the Lagrange multiplier $\hat{\lambda}$ of ETEL.

$$\begin{bmatrix} & \hat{\tau}^{ETEL} \\ \text{Auxiliary parameter of ETEL } \rightarrow & \hat{\kappa}^{ETEL} \\ \text{Lagrange multiplier of ETEL } \rightarrow & \hat{\lambda}^{ETEL} \\ & \hat{\theta}^{ETEL} \end{bmatrix} - \begin{bmatrix} \hat{\tau}^{EL} & & \\ \hat{\kappa}^{EL} & \leftarrow & \text{Lagrange multiplier of EL} \\ \hat{\lambda}^{EL} & \leftarrow & \text{parameter added for convenience} \\ \hat{\theta}^{EL} & & \end{bmatrix}$$

Hence, our results do not imply that the Lagrange multiplier EL and ETEL are equivalent up to $O_p\left(n^{-1}\right)$ but rather that the auxiliary parameter $\hat{\kappa}^{ETEL}$ in fact plays the role of EL's Lagrange multiplier in the ETEL estimator. Since the influence function of EL and ETEL are the same, we have that

$$\bar{Q}_l^{ETEL} - \bar{Q}_l^{EL} = \left(\bar{\Psi}_{l,j}^{ETEL} - \bar{\Psi}_{l,j}^{EL}\right)\bar{\Psi}_j + \bar{\Psi}_j\left(\Psi_{l,jk}^{ETEL} - \Psi_{l,jk}^{EL}\right)\bar{\Psi}_k/2$$

5.1.1 Calculation of $\left(\bar{\Psi}_{l,j}^{ETEL} - \bar{\Psi}_{l,j}^{EL}\right)\bar{\Psi}_j$

$\left(\bar{\Psi}_{l,j}^{ETEL} - \bar{\Psi}_{l,j}^{EL}\right)\bar{\Psi}_j = \Phi_{lh}^{-1}\left(\bar{\Phi}_{h,j}^{ETEL} - \bar{\Phi}_{h,j}^{EL}\right)\bar{\Psi}_j$, where

$$\left(\bar{\Phi}_{\cdot,\cdot}^{ETEL} - \bar{\Phi}_{\cdot,\cdot}^{EL}\right)\bar{\Psi}_{\cdot} = \left(\begin{bmatrix} -1 & 0 & \bar{g}' & 0 \\ 0 & 0 & \bar{g}\bar{g}' & \bar{G} \\ \bar{g} & \bar{g}\bar{g}' & -\bar{g}\bar{g}' & 0 \\ 0 & \bar{G}' & 0 & 0 \end{bmatrix} - \begin{bmatrix} -1 & 0 & \bar{g}' & 0 \\ 0 & 0 & \bar{g}\bar{g}' & \bar{G} \\ 0 & \bar{g}\bar{g}' & -\bar{g}\bar{g}' & 0 \\ 0 & \bar{G}' & 0 & 0 \end{bmatrix}\right)\begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} =$$

$$= \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ \bar{g} & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}\begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \\ 0 \end{bmatrix}$$

Thus, $\left(\bar{\Psi}_{\cdot,\cdot}^{ETEL} - \bar{\Psi}_{\cdot,\cdot}^{EL}\right)\bar{\Psi}_{\cdot} = \mathbf{0}$.

5.1.2 Calculation of $\bar{\Psi}_j\left(\Psi_{l,jk}^{ETEL} - \Psi_{l,jk}^{EL}\right)\bar{\Psi}_k/2$

$\bar{\Psi}_j\left(\Psi_{l,jk}^{ETEL} - \Psi_{l,jk}^{EL}\right)\bar{\Psi}_k = \Phi_{lh}^{-1}\bar{\Psi}_j\left(\Phi_{h,jk}^{ETEL} - \Phi_{h,jk}^{EL}\right)\bar{\Psi}_k = \Phi_{lh}^{-1}\bar{\Psi}_{\cdot}' E^*\left[\frac{\partial^2}{\partial\beta\partial\beta'}\left(\phi_h^{ETEL} - \phi_h^{EL}\right)\right]\bar{\Psi}_{\cdot}$
where

$$\left(\dot{\phi}^{ETEL} - \dot{\phi}^{EL}\right) = \left(\begin{bmatrix} \dot{\tau} - \tau \\ \dot{\tau}\dot{g} \\ (\tau - \dot{\tau})\dot{g} + \dot{\tau}\dot{g}\dot{g}'\kappa \\ \dot{\tau}\dot{G}'\kappa + \dot{\tau}\dot{G}'\lambda\dot{g}'\kappa - \dot{\tau}\dot{G}'\lambda + \tau\dot{G}'\lambda \end{bmatrix} - \begin{bmatrix} \dot{\tau} - \tau \\ \dot{\tau}\dot{g} \\ \dot{\varepsilon}\dot{g} - \dot{\tau}\dot{g} \\ \dot{\varepsilon}\dot{G}'\kappa \end{bmatrix}\right)$$

$$= \begin{bmatrix} 0 \\ 0 \\ (\tau - 1)\dot{g} + (\dot{\tau} - \dot{\varepsilon})\dot{g}\dot{g}'\kappa \\ (\dot{\tau} - \dot{\varepsilon})\dot{G}'\kappa + \dot{\tau}\dot{G}'\lambda\dot{g}'\kappa - (\dot{\tau} - \tau)\dot{G}'\lambda \end{bmatrix}$$

where we used the identity $\dot{\varepsilon}\dot{g} = \frac{\dot{g}}{1-\dot{g}'\kappa} = \dot{g}\left(1 + \frac{\dot{g}'\kappa}{1-\dot{g}'\kappa}\right) = \dot{g}\left(1 + \dot{\varepsilon}\dot{g}'\kappa\right) = \dot{g} + \dot{\varepsilon}\dot{g}\dot{g}'\kappa$.
To calculate $\left(\Phi_{h,jk}^{ETEL} - \Phi_{h,jk}^{EL}\right)$:

$$\left[\frac{\partial^2}{\partial\beta\partial\beta'}\left(\dot{\phi}_{l_\lambda+h}^{ETEL} - \dot{\phi}_{l_\lambda+h}^{EL}\right)\right]_{\beta=\beta^*} = \left[\frac{\partial^2}{\partial\beta\partial\beta'}\left((\tau - 1)\dot{g}_h + (\dot{\tau} - \dot{\varepsilon})\dot{g}_h\dot{g}'\kappa\right)\right]_{\beta=\beta^*} =$$

$$= \begin{bmatrix} 0 & 0 & 0 & \dot{G}_{h\cdot} \\ 0 & -2\dot{g}\dot{g}'\dot{g}_h & \dot{g}_h\dot{g}\dot{g}' & 0 \\ 0 & \dot{g}_h\dot{g}\dot{g}' & 0 & 0 \\ \dot{G}_{h\cdot}' & 0 & 0 & 0 \end{bmatrix}_{\beta=\beta^*}$$

and

$$\left[\frac{\partial^2}{\partial\beta\partial\beta'}\left(\dot{\phi}_{l_\theta+h}^{ETEL} - \dot{\phi}_{l_\theta+h}^{EL}\right)\right]_{\beta=\beta^*} = \left[\frac{\partial^2}{\partial\beta\partial\beta'}\left((\dot{\tau} - \dot{\varepsilon})\dot{G}_{\cdot h}'\kappa + \dot{\tau}\dot{G}_{\cdot h}'\lambda\dot{g}'\kappa + (\tau - \dot{\tau})\dot{G}_{\cdot h}'\lambda\right)\right]_{\beta=\beta^*}$$

$$= \begin{bmatrix} 0 & 0 & \dot{G}'_{\cdot h} & 0 \\ 0 & -\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h} & \frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h} & 0 \\ \dot{G}_{\cdot h} & \frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h} & -\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h} & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}_{\beta=\beta^*}$$

Then

$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\tau+h,jk} - \Phi^{EL}_{l_\tau+h,jk}\right)\bar{\Psi}_k = 0$

$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\kappa+h,jk} - \Phi^{EL}_{l_\kappa+h,jk}\right)\bar{\Psi}_k = 0$

$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\lambda+h,jk} - \Phi^{EL}_{l_\lambda+h,jk}\right)\bar{\Psi}_k =$

$$= \begin{bmatrix} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & G_{h\cdot} \\ 0 & -2E^*\left[\dot{g}\dot{g}'\dot{g}_h\right] & E^*\left[\dot{g}_h\dot{g}\dot{g}'\right] & 0 \\ 0 & E^*\left[\dot{g}_h\dot{g}\dot{g}'\right] & 0 & 0 \\ G'_{h\cdot} & 0 & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} = 0$$

where $G'_{\cdot h} \equiv \left(G_{\cdot h}\right)'$.

$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\theta+h,jk} - \Phi^{EL}_{l_\theta+h,jk}\right)\bar{\Psi}_k =$

$$= \begin{bmatrix} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{bmatrix} \begin{matrix} 0 & 0 & G'_{\cdot h} & 0 \\ 0 & -E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right] & E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right] & 0 \\ G_{\cdot h} & E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right] & -E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right] & 0 \\ 0 & 0 & 0 & 0 \end{matrix} \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix}$$

$= 0$

Since $\bar{\Psi}_j\left(\Phi^{ETEL}_{h,jk} - \Phi^{EL}_{h,jk}\right)\bar{\Psi}_k = 0$ for all h, it follows that $\bar{\Psi}_j\left(\Psi^{ETEL}_{l,jk} - \Psi^{EL}_{l,jk}\right)\bar{\Psi}_k = 0$ for all l.

Conclusion: $\bar{Q}^{ETEL}_l - \bar{Q}^{EL}_l = 0$.

5.2 Calculation of $\bar{Q}^{ETEL}_l = \bar{Q}^{EL}_l \equiv \bar{Q}_l$

$$\bar{Q}^{EL}_l = \bar{\Psi}^{EL}_{l,j}\bar{\Psi}_j + \bar{\Psi}_j\Psi^{EL}_{l,jk}\bar{\Psi}_k/2$$

5.2.1 Calculation of $\bar{\Psi}^{EL}_{l,j}\bar{\Psi}_j$

$$\bar{\Psi}^{EL}_{\cdot,\cdot} = -\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & P & P & H' \\ 0 & P & \left(P-\Omega^{-1}\right) & H' \\ 0 & H & H & -\Sigma \end{bmatrix} \begin{bmatrix} 0 & 0 & \bar{g}' & 0 \\ 0 & 0 & \bar{\Omega} & \bar{G} \\ 0 & \bar{\Omega} & -\bar{\Omega} & 0 \\ 0 & \bar{G}' & 0 & 0 \end{bmatrix} = -\begin{bmatrix} 0 & 0 & -\bar{g}' & 0 \\ 0 & P\bar{\Omega}+H'\bar{G}' & 0 & P\bar{G} \\ 0 & \left(P-\Omega^{-1}\right)\bar{\Omega}+H'\bar{G}' & \Omega^{-1}\bar{\Omega} & P\bar{G} \\ 0 & H\bar{\Omega}-\Sigma\bar{G}' & 0 & H\bar{G} \end{bmatrix}$$

$$\bar{\Psi}^{EL}_{\cdot,\cdot}\bar{\Psi}_{\cdot} = -\begin{bmatrix} 0 & 0 & -\bar{g}' & 0 \\ 0 & P\bar{\Omega}+H'\bar{G}' & 0 & P\bar{G} \\ 0 & P\bar{\Omega}-\Omega^{-1}\bar{\Omega}+H'\bar{G}' & \Omega^{-1}\bar{\Omega} & P\bar{G} \\ 0 & H\bar{\Omega}-\Sigma\bar{G}' & 0 & H\bar{G} \end{bmatrix} \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} = \begin{bmatrix} -\bar{g}'P\bar{g} \\ P\bar{\Omega}P\bar{g}+H'\bar{G}'P\bar{g}+P\bar{G}H\bar{g} \\ P\bar{\Omega}P\bar{g}+H'\bar{G}'P\bar{g}+P\bar{G}H\bar{g} \\ H\bar{\Omega}P\bar{g}-\Sigma\bar{G}'P\bar{g}+H\bar{G}H\bar{g} \end{bmatrix}$$

5.2.2 Calculation of $\bar{\Psi}_j\Psi^{EL}_{l,jk}\bar{\Psi}_k/2$

$\bar{\Psi}_j\Psi^{EL}_{l,jk}\bar{\Psi}_k/2 = -\Phi^{-1}_{lh}\bar{\Psi}_j\Phi^{EL}_{h,jk}\bar{\Psi}_k/2$ where

$\bar{\Psi}_j\Phi^{EL}_{l_\tau+h,jk}\bar{\Psi}_k = \bar{\Psi}_jE^*\left[\frac{\partial^2}{\partial\beta_j\partial\beta_k}\dot{\phi}^{EL}_{l_\tau+h}\right]\bar{\Psi}_k = \bar{\Psi}'_{\cdot}E^*\left[\frac{\partial^2\left(\dot{\tau}-\tau\right)}{\partial\beta\partial\beta'}\right]\bar{\Psi}_{\cdot} =$

$$= \begin{bmatrix} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & \Omega & G \\ 0 & 0 & G' & 0 \end{bmatrix} \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} = g'\left(P\Omega P\right)g + g'H'\left(G'P\right)g + g'\left(PG\right)Hg$$

$= \bar{g}'P\bar{g} + 0 + 0$

$\bar{\Psi}_j\Phi^{EL}_{l_\kappa+h,jk}\bar{\Psi}_k = \bar{\Psi}_jE^*\left[\frac{\partial^2}{\partial\beta_j\partial\beta_k}\dot{\phi}^{EL}_{l_\kappa+h}\right]\bar{\Psi}_k = \bar{\Psi}'_{\cdot}E^*\left[\frac{\partial^2\left(\dot{\tau}\dot{g}_h\right)}{\partial\beta\partial\beta'}\right]\bar{\Psi}_{\cdot} =$

$$= \begin{bmatrix} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 \\ 0 & 0 & E^*\left[\dot{g}_h\dot{g}\dot{g}'\right] & E^*\left[\frac{\partial}{\partial\theta'}\left(\dot{g}_h\dot{g}\right)\right] \\ 0 & 0 & E^*\left[\frac{\partial}{\partial\theta}\left(\dot{g}_h\dot{g}'\right)\right] & E^*\left[\frac{\partial^2\dot{g}_h}{\partial\theta\partial\theta'}\right] \end{bmatrix} \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} =$$

$$= \bar{g}'PE^*\left[\dot{g}_h\dot{g}\dot{g}'\right]P\bar{g} + \bar{g}'PE^*\left[\frac{\partial}{\partial\theta'}\left(\dot{g}_h\dot{g}\right)\right]H\bar{g} + \bar{g}'H'E^*\left[\frac{\partial}{\partial\theta}\left(\dot{g}_h\dot{g}'\right)\right]P\bar{g} + \bar{g}'H'E^*\left[\frac{\partial^2\dot{g}_h}{\partial\theta\partial\theta'}\right]H\bar{g}$$

$$= \left(\left(\bar{g}'P\right)_j E^*\left[\dot{g}\dot{g}_j\dot{g}'\right]P\bar{g}\right)_h + \left(\left(\bar{g}'P\right)_j E^*\left[\frac{\partial}{\partial\theta'}\left(\dot{g}\dot{g}_j\right)\right]H\bar{g}\right)_h + \left(\left(\bar{g}'H'\right)_j E^*\left[\frac{\partial}{\partial\theta_j}\left(\dot{g}\dot{g}'\right)\right]P\bar{g}\right)_h +$$
$$+ \left(\left(\bar{g}'H'\right)_j E^*\left[\frac{\partial^2\dot{g}}{\partial\theta_j\partial\theta'}\right]H\bar{g}\right)_h$$

$$\bar{\Psi}_j\Phi^{EL}_{l_\lambda+h,jk}\bar{\Psi}_k = \bar{\Psi}_j E^*\left[\frac{\partial^2}{\partial\beta_j\partial\beta_k}\dot{\phi}^{EL}_{l_\lambda+h}\right]_{\beta=\beta^*}\bar{\Psi}_k = \bar{\Psi}'_{\cdot}E^*\left[\frac{\partial^2\left(\dot{\varepsilon}\dot{g}_h - \dot{\tau}\dot{g}_h\right)}{\partial\beta\partial\beta'}\right]_{\beta=\beta^*}\bar{\Psi}_{\cdot} =$$

$$= \begin{bmatrix} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 2E^*\left[\dot{g}\dot{g}'\dot{g}_h\right] & 0 & E^*\left[\frac{\partial\left(\dot{g}\dot{g}_h\right)}{\partial\theta'}\right] \\ 0 & 0 & -E^*\left[\dot{g}\dot{g}'\dot{g}_h\right] & -E^*\left[\frac{\partial\left(\dot{g}\dot{g}_h\right)}{\partial\theta'}\right] \\ 0 & E^*\left[\frac{\partial\left(\dot{g}_h\dot{g}'\right)}{\partial\theta}\right] & -E^*\left[\frac{\partial\left(\dot{g}_h\dot{g}'\right)}{\partial\theta}\right] & 0 \end{bmatrix} \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} =$$

$$= \bar{g}'PE^*\left[\dot{g}\dot{g}'\dot{g}_h\right]P\bar{g} = \left(\left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\right)_h$$

$$\bar{\Psi}_j\Phi^{EL}_{l_\theta+h,jk}\bar{\Psi}_k = \bar{\Psi}_j E^*\left[\frac{\partial^2}{\partial\beta_j\partial\beta_k}\dot{\phi}^{EL}_{l_\theta+h}\right]_{\beta=\beta^*}\bar{\Psi}_k = \bar{\Psi}'_{\cdot}E^*\left[\frac{\partial^2\left(\dot{\varepsilon}\dot{G}'_{\cdot h}\kappa\right)}{\partial\beta\partial\beta'}\right]_{\beta=\beta^*}\bar{\Psi}_{\cdot} =$$

$$= \begin{bmatrix} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{bmatrix} \begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right] & 0 & E^*\left[\frac{\partial^2\dot{g}}{\partial\theta_h\partial\theta'}\right] \\ 0 & 0 & 0 & 0 \\ 0 & E^*\left[\frac{\partial^2\dot{g}'}{\partial\theta\partial\theta_h}\right] & 0 & 0 \end{bmatrix} \begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix} =$$

$$= \bar{g}'PE^*\left[\frac{\partial\left(\dot{g}_j\dot{g}'\right)}{\partial\theta_h}\right]P\bar{g} + \bar{g}'PE^*\left[\frac{\partial^2\dot{g}}{\partial\theta_h\partial\theta'}\right]H\bar{g} + \bar{g}'H'E^*\left[\frac{\partial^2\dot{g}'}{\partial\theta\partial\theta_h}\right]P\bar{g}$$

$$= \left(\left(\bar{g}'P\right)_j E^*\left[\frac{\partial\left(\dot{g}_j\dot{g}'\right)}{\partial\theta}\right]P\bar{g}\right)_h + \left(\left(\bar{g}'P\right)_j E^*\left[\frac{\partial^2\dot{g}_j}{\partial\theta\partial\theta'}\right]H\bar{g}\right)_h + \left(\left(\bar{g}'H'\right)_j E^*\left[\frac{\partial^2\dot{g}'}{\partial\theta\partial\theta_j}\right]P\bar{g}\right)_h$$

Collecting these 4 results into a single vector, we obtain:

$$\bar{\Psi}_j\Phi^{EL}_{\cdot,jk}\bar{\Psi}_k = \begin{bmatrix} \bar{g}'P\bar{g} \\ \left(\bar{g}'P\right)_j E^*\left[\dot{g}\dot{g}_j\dot{g}'\right]P\bar{g} + \left(\bar{g}'P\right)_j E^*\left[\frac{\partial\left(\dot{g}\dot{g}_j\right)}{\partial\theta'}\right]H\bar{g} + \left(\bar{g}'H'\right)_j E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_j}\right]P\bar{g} + \left(\bar{g}'H'\right)_j E^*\left[\frac{\partial^2\dot{g}}{\partial\theta_j\partial\theta'}\right]H\bar{g} \\ \left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g} \\ \left(\bar{g}'P\right)_j E^*\left[\frac{\partial\left(\dot{g}_j\dot{g}'\right)}{\partial\theta}\right]P\bar{g} + \left(\bar{g}'P\right)_j E^*\left[\frac{\partial^2\dot{g}_j}{\partial\theta\partial\theta'}\right]H\bar{g} + \left(\bar{g}'H'\right)_j E^*\left[\frac{\partial^2\dot{g}'}{\partial\theta\partial\theta_j}\right]P\bar{g} \end{bmatrix}$$

$$\bar{\Psi}_j\Psi^{EL}_{\cdot,jk}\bar{\Psi}_k = -\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & P & P & H' \\ 0 & P & \left(P-\Omega^{-1}\right) & H' \\ 0 & H & H & -\Sigma \end{bmatrix} \times$$
$$\times \begin{bmatrix} \bar{g}'P\bar{g} \\ \left(\bar{g}'P\right)_j E^*\left[\dot{g}\dot{g}_j\dot{g}'\right]P\bar{g} + \left(\bar{g}'P\right)_j E^*\left[\frac{\partial}{\partial\theta'}\left(\dot{g}\dot{g}_j\right)\right]H\bar{g} + \left(\bar{g}'H'\right)_j E^*\left[\frac{\partial}{\partial\theta_j}\left(\dot{g}\dot{g}'\right)\right]P\bar{g} + \left(\bar{g}'H'\right)_j E^*\left[\frac{\partial^2\dot{g}}{\partial\theta_j\partial\theta'}\right]H\bar{g} \\ \left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g} \\ \left(\bar{g}'P\right)_j E^*\left[\frac{\partial\left(\dot{g}_j\dot{g}'\right)}{\partial\theta}\right]P\bar{g} + \left(\bar{g}'P\right)_j E^*\left[\frac{\partial^2\dot{g}_j}{\partial\theta\partial\theta'}\right]H\bar{g} + \left(\bar{g}'H'\right)_j E^*\left[\frac{\partial^2\dot{g}'}{\partial\theta\partial\theta_j}\right]P\bar{g} \end{bmatrix}$$
$$= \begin{bmatrix} \bar{g}'P\bar{g} \\ \Xi_1 \\ \Xi_1 + \Omega^{-1}\left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g} \\ \Xi_2 \end{bmatrix}$$

where

$$\Xi_1 = -P(\left(\bar{g}'P\right)_j E^*\left[\dot{g}\dot{g}_j\dot{g}'\right]P\bar{g} + \left(\bar{g}'P\right)_j E^*\left[\frac{\partial}{\partial\theta'}\left(\dot{g}\dot{g}_j\right)\right]H\bar{g} + \left(\bar{g}'H'\right)_j E^*\left[\frac{\partial}{\partial\theta_j}\left(\dot{g}\dot{g}'\right)\right]P\bar{g} +$$

$$+\left(\bar{g}'H'\right)_j E^*\left[\tfrac{\partial^2\dot{g}}{\partial\theta_j\partial\theta'}\right]H\bar{g}+\left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g})+$$

$$-H'\left(\left(\bar{g}'P\right)_j E^*\left[\frac{\partial\left(\dot{g}_j\dot{g}'\right)}{\partial\theta}\right]P\bar{g}+\left(\bar{g}'P\right)_j E^*\left[\tfrac{\partial^2\dot{g}_j}{\partial\theta\partial\theta'}\right]H\bar{g}+\left(\bar{g}'H'\right)_j E^*\left[\tfrac{\partial^2\dot{g}'}{\partial\theta\partial\theta_j}\right]P\bar{g}\right)$$

$$\Xi_2=-H((\bar{g}'P)_j E^*\left[\dot{g}\dot{g}_j\dot{g}'\right]P\bar{g}+\left(\bar{g}'P\right)_j E^*\left[\tfrac{\partial}{\partial\theta'}\left(\dot{g}\dot{g}_j\right)\right]H\bar{g}+\left(\bar{g}'H'\right)_j E^*\left[\tfrac{\partial}{\partial\theta_j}\left(\dot{g}\dot{g}'\right)\right]P\bar{g}+$$

$$+\left(\bar{g}'H'\right)_j E^*\left[\tfrac{\partial^2\dot{g}}{\partial\theta_j\partial\theta'}\right]H\bar{g}+\left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g})+$$

$$+\Sigma\left(\left(\bar{g}'P\right)_j E^*\left[\frac{\partial\left(\dot{g}_j\dot{g}'\right)}{\partial\theta}\right]P\bar{g}+\left(\bar{g}'P\right)_j E^*\left[\tfrac{\partial^2\dot{g}_j}{\partial\theta\partial\theta'}\right]H\bar{g}+\left(\bar{g}'H'\right)_j E^*\left[\tfrac{\partial^2\dot{g}'}{\partial\theta\partial\theta_j}\right]P\bar{g}\right)$$

Finally,

$$\bar{Q}^{EL}_{\cdot}=\bar{\Psi}^{EL}_{\cdot,j}\bar{\Psi}_j+\bar{\Psi}_j\Psi^{EL}_{\cdot,jk}\bar{\Psi}_k/2=\left[\begin{array}{c}-\bar{g}'P\bar{g}\\ P\bar{\Omega}P\bar{g}+H'\bar{G}'P\bar{g}+P\bar{G}H\bar{g}\\ P\bar{\Omega}P\bar{g}+H'\bar{G}'P\bar{g}+P\bar{G}H\bar{g}\\ H\bar{\Omega}P\bar{g}-\Sigma\bar{G}'P\bar{g}+H\bar{G}H\bar{g}\end{array}\right]+\tfrac{1}{2}\left[\begin{array}{c}\bar{g}'P\bar{g}\\ \Xi_1\\ \Xi_1+\Omega^{-1}\left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\\ \Xi_2\end{array}\right]$$

$$=\left[\begin{array}{c}-\bar{g}'P\bar{g}/2\\ \Xi_3\\ \Xi_3+\tfrac{1}{2}\Omega^{-1}\left(\bar{g}'P\right)_j E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\\ \Xi_4\end{array}\right]$$

where

$$\Xi_3=\tfrac{1}{2}\Xi_1+P\bar{\Omega}P\bar{g}+H'\bar{G}'P\bar{g}+P\bar{G}H\bar{g}$$
$$\Xi_4=\tfrac{1}{2}\Xi_2+H\bar{\Omega}P\bar{g}-\Sigma\bar{G}'P\bar{g}+H\bar{G}H\bar{g}$$

6 Calculation of $\bar{R}$.

We will prove that $E\left[\left(\bar{R}^{ETEL}_{l_\theta+l}-\bar{R}^{EL}_{l_\theta+l}\right)\bar{\Psi}_{l_\theta+m}\right]=o\left(n^{-2}\right)$. This will be done by showing that most of the term comprised in $\bar{R}^{ETEL}_{l_\theta+l}-\bar{R}^{EL}_{l_\theta+l}$ are zero. The remaining nonzero terms will be shown to be uncorrelated with $\bar{\Psi}_{l_\theta+m}$ (up to $O\left(n^{-2}\right)$).
Since $\bar{Q}^{ETEL}_j=\bar{Q}^{EL}_j\equiv\bar{Q}_j$

$$\begin{aligned}R^{ETEL}_l-R^{EL}_l &= \left(\bar{\Psi}^{ETEL}_{l,j}-\bar{\Psi}^{EL}_{l,j}\right)\bar{Q}_j+\left(\Psi^{ETEL}_{l,jk}-\Psi^{EL}_{l,jk}\right)\bar{\Psi}_k\bar{Q}_j+\\ &\quad+\frac{1}{2}\left(\bar{\Psi}^{ETEL}_{l,jk}-\bar{\Psi}^{EL}_{l,jk}\right)\bar{\Psi}_j\bar{\Psi}_k+\frac{1}{6}\left(\Psi^{ETEL}_{l,jkh}-\Psi^{EL}_{l,jkh}\right)\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_h\end{aligned}$$

6.1 Calculation of $\left(\bar{\Psi}^{ETEL}_{l_\theta+l,j}-\bar{\Psi}^{EL}_{l_\theta+l,j}\right)\bar{Q}_j$

$$\left(\bar{\Psi}^{ETEL}_{l_\theta+l,\cdot}-\bar{\Psi}^{EL}_{l_\theta+l,\cdot}\right)=\Phi_{l_\theta+l,h}\left(\bar{\Phi}^{ETEL}_{h,\cdot}-\bar{\Phi}^{EL}_{h,\cdot}\right)=$$

$$=-\left[\begin{array}{cccc}0 & H_{l\cdot} & H_{l\cdot} & -\Sigma_{l\cdot}\end{array}\right]\left(\left[\begin{array}{cccc}-1 & 0 & \bar{g}' & 0\\ 0 & 0 & \bar{g}\bar{g}' & \bar{G}\\ \bar{g} & \bar{g}\bar{g}' & -\bar{g}\bar{g}' & 0\\ 0 & \bar{G}' & 0 & 0\end{array}\right]-\left[\begin{array}{cccc}-1 & 0 & \bar{g}' & 0\\ 0 & 0 & \bar{g}\bar{g}' & \bar{G}\\ 0 & \bar{g}\bar{g}' & -\bar{g}\bar{g}' & 0\\ 0 & \bar{G}' & 0 & 0\end{array}\right]\right)=$$

$$=\left[\begin{array}{cccc}-H_{l\cdot}\bar{g} & 0 & 0 & 0\end{array}\right]$$

$$\left(\bar{\Psi}^{ETEL}_{l_\theta+l,j}-\bar{\Psi}^{EL}_{l_\theta+l,j}\right)\bar{Q}_j=\left(\bar{\Psi}^{ETEL}_{l_\theta+l,1}-\bar{\Psi}^{EL}_{l_\theta+l,1}\right)\bar{Q}_1=-H_{l\cdot}\bar{g}\left(-\bar{g}'P\bar{g}/2\right)=H_{l\cdot}\bar{g}\bar{g}'P\bar{g}/2$$

6.2 Calculation of $\left(\Psi^{ETEL}_{l,jk}-\Psi^{EL}_{l,jk}\right)\bar{\Psi}_k\bar{Q}_j$

$\left(\Psi^{ETEL}_{l,jk}-\Psi^{EL}_{l,jk}\right)\bar{\Psi}_k\bar{Q}_j=\Phi^{-1}_{lh}\left(\Phi^{ETEL}_{h,jk}-\Phi^{EL}_{h,jk}\right)\bar{\Psi}_k\bar{Q}_j$ where $\left(\Phi^{ETEL}_{h,jk}-\Phi^{EL}_{h,jk}\right)\bar{\Psi}_k$ can be calculated as in Section 5.1.2:

Since $\left(\dot{\phi}^{ETEL}-\dot{\phi}^{EL}\right)=\left[\begin{array}{c}0\\ 0\\ \left(\tau-1\right)\dot{g}+\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{g}\dot{g}'\kappa\\ \left(\dot{\tau}-\dot{\varepsilon}\right)\dot{G}'\kappa+\dot{\tau}\dot{G}'\lambda\dot{g}'\kappa-\left(\dot{\tau}-\tau\right)\dot{G}'\lambda\end{array}\right]$

$$\left(\Phi^{ETEL}_{l_\tau+h,\cdot k}-\Phi^{EL}_{l_\tau+h,\cdot k}\right)\bar{\Psi}_k=\mathbf{0}$$

$$\left(\Phi^{ETEL}_{l_\kappa+h,\cdot k}-\Phi^{EL}_{l_\kappa+h,\cdot k}\right)\bar{\Psi}_k=\mathbf{0}$$

$$\left(\Phi^{ETEL}_{l_\lambda+h,\cdot k}-\Phi^{EL}_{l_\lambda+h,\cdot k}\right)\bar{\Psi}_k=\begin{bmatrix}0&0&0&G_{h\cdot}\\0&-2E^*\left[\dot{g}\dot{g}'\dot{g}_h\right]&E^*\left[\dot{g}_h\dot{g}\dot{g}'\right]&0\\0&E^*\left[\dot{g}_h\dot{g}\dot{g}'\right]&0&0\\G'_{h\cdot}&0&0&0\end{bmatrix}\begin{bmatrix}0\\-P\bar{g}\\-P\bar{g}\\-H\bar{g}\end{bmatrix}=\begin{bmatrix}-G_{h\cdot}H\bar{g}\\E^*\left[\dot{g}\dot{g}'\dot{g}_h\right]P\bar{g}\\-E^*\left[\dot{g}\dot{g}'\dot{g}_h\right]P\bar{g}\\0\end{bmatrix}$$

$$\left(\Phi^{ETEL}_{l_\theta+h,\cdot k}-\Phi^{EL}_{l_\theta+h,\cdot k}\right)\bar{\Psi}_k=\begin{bmatrix}0&0&G'_{\cdot h}&0\\0&-E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right]&E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right]&0\\G_{\cdot h}&E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right]&-E^*\left[\frac{\partial\left(\dot{g}\dot{g}'\right)}{\partial\theta_h}\right]&0\\0&0&0&0\end{bmatrix}\begin{bmatrix}0\\-P\bar{g}\\-P\bar{g}\\-H\bar{g}\end{bmatrix}=\begin{bmatrix}-G'_{\cdot h}P\bar{g}\\0\\0\\0\end{bmatrix}=\begin{bmatrix}0\\0\\0\\0\end{bmatrix}$$

Then, since $\left(\Phi^{ETEL}_{h,jk}-\Phi^{EL}_{h,jk}\right)\bar{\Psi}_k\bar{Q}_j=\bar{Q}'_{\cdot}\left(\Phi^{ETEL}_{h,\cdot k}-\Phi^{EL}_{h,\cdot k}\right)\bar{\Psi}_k$, we obtain

$$\bar{Q}'_{\cdot}\left(\Phi^{ETEL}_{l_\tau+h,\cdot k}-\Phi^{EL}_{l_\tau+h,\cdot k}\right)\bar{\Psi}_k=0$$

$$\bar{Q}'_{\cdot}\left(\Phi^{ETEL}_{l_\kappa+h,\cdot k}-\Phi^{EL}_{l_\kappa+h,\cdot k}\right)\bar{\Psi}_k=0$$

$$\bar{Q}'_{\cdot}\left(\Phi^{ETEL}_{l_\lambda+h,\cdot k}-\Phi^{EL}_{l_\lambda+h,\cdot k}\right)\bar{\Psi}_k=\begin{bmatrix}-\bar{g}'P\bar{g}/2&\Xi'_3&\Xi'_3+\frac{1}{2}\left(\bar{g}'P\right)_j\bar{g}'PE^*\left[\dot{g}_j\dot{g}\dot{g}'\right]\Omega^{-1}&\Xi'_4\end{bmatrix}\begin{bmatrix}-G_{h\cdot}H\bar{g}\\E^*\left[\dot{g}\dot{g}'\dot{g}_h\right]P\bar{g}\\-E^*\left[\dot{g}\dot{g}'\dot{g}_h\right]P\bar{g}\\0\end{bmatrix}$$

$$=\tfrac{1}{2}\bar{g}'P\bar{g}G_{h\cdot}H\bar{g}-\tfrac{1}{2}\left(\bar{g}'P\right)_j\bar{g}'PE^*\left[\dot{g}_j\dot{g}\dot{g}'\right]\Omega^{-1}E^*\left[\dot{g}\dot{g}'\dot{g}_h\right]P\bar{g}=$$

$$=\tfrac{1}{2}G_{h\cdot}H\bar{g}\bar{g}'P\bar{g}-\tfrac{1}{2}E^*\left[\dot{g}_h\dot{g}_k\dot{g}'\right]\Omega^{-1}E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\left(\bar{g}'P\right)_j\left(\bar{g}'P\right)_k$$

$$\bar{Q}'_{\cdot}\left(\Phi^{ETEL}_{l_\theta+h,\cdot k}-\Phi^{EL}_{l_\theta+h,\cdot k}\right)\bar{\Psi}_k=0$$

Collecting the previous four quantities into a single vector, we have

$$\left(\Phi^{ETEL}_{\cdot,jk}-\Phi^{EL}_{\cdot,\cdot jk}\right)\bar{Q}_j\bar{\Psi}_k=\begin{bmatrix}0\\0\\\frac{1}{2}GH\bar{g}\bar{g}'P\bar{g}+E^*\left[\dot{g}\dot{g}_k\dot{g}'\right]\Omega^{-1}E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\left(\bar{g}'P\right)_j\left(\bar{g}'P\right)_k\\0\end{bmatrix}$$

$$\Phi^{-1}_{l_\theta+l,\cdot}\left(\Phi^{ETEL}_{\cdot,jk}-\Phi^{EL}_{\cdot,\cdot jk}\right)\bar{Q}_j\bar{\Psi}_k=$$

$$=-\begin{bmatrix}0&H_{l\cdot}&H_{l\cdot}&-\Sigma_{l\cdot}\end{bmatrix}\begin{bmatrix}0\\0\\\frac{1}{2}GH\bar{g}\bar{g}'P\bar{g}+E^*\left[\dot{g}\dot{g}_k\dot{g}'\right]\Omega^{-1}E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\left(\bar{g}'P\right)_j\left(\bar{g}'P\right)_k\\0\end{bmatrix}$$

$$=-\left(\tfrac{1}{2}HGH\bar{g}\bar{g}'P\bar{g}\right)_l-\left(HE^*\left[\dot{g}\dot{g}_k\dot{g}'\right]\Omega^{-1}E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\left(\bar{g}'P\right)_j\left(\bar{g}'P\right)_k\right)_l$$

$$=-\tfrac{1}{2}\left(\left(\left(G'\Omega^{-1}G\right)^{-1}G'\Omega^{-1}\right)GH\bar{g}\bar{g}'P\bar{g}\right)_l-\left(HE^*\left[\dot{g}\dot{g}_k\dot{g}'\right]\Omega^{-1}E^*\left[\dot{g}_j\dot{g}\dot{g}'\right]P\bar{g}\left(\bar{g}'P\right)_j\left(\bar{g}'P\right)_k\right)_l$$

$$=-\left(\tfrac{1}{2}H\bar{g}\bar{g}'P\bar{g}\right)_l-\left(HE^*\left[\dot{g}\dot{g}_k\dot{g}'\right]\Omega^{-1}E^*\left[\dot{g}_j\dot{g}\dot{g}_m\right]\left(P\bar{g}\right)_m\left(P\bar{g}\right)_j\left(P\bar{g}\right)_k\right)_l$$

The first term is exactly cancels the first term of the expansion of $\bar{R}$. (see section 6.1).
The second term is uncorrelated with $\bar{\Psi}_{l_\theta+l}$, which implies that it does not contribute to the n^{-2} variance. Indeed, all the random contributions to the second term are of the form $P\bar{g}$, while $\bar{\Psi}_{l_\theta+l}$ is of the form $H\bar{g}$ and $E^*\left[P\bar{g}\bar{g}'H\right]=P\Omega H=0$.

6.3 Calculation of $\bar{\Psi}_j\left(\bar{\Psi}^{ETEL}_{l,jk}-\bar{\Psi}^{EL}_{l,jk}\right)\bar{\Psi}_k$

$$\bar{\Psi}_j\left(\bar{\Psi}^{ETEL}_{l,jk}-\bar{\Psi}^{EL}_{l,jk}\right)\bar{\Psi}_k=\Phi^{-1}_{lh}\bar{\Psi}_j\left(\bar{\Phi}^{ETEL}_{h,jk}-\bar{\Phi}^{EL}_{h,jk}\right)\bar{\Psi}_k$$

Since $\left(\dot{\phi}^{ETEL}-\dot{\phi}^{EL}\right)=\begin{bmatrix}0\\0\\\left(\tau-1\right)\dot{g}+\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{g}\dot{g}'\kappa\\\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{G}'\kappa+\dot{\tau}\dot{G}'\lambda\dot{g}'\kappa-\left(\dot{\tau}-\tau\right)\dot{G}'\lambda\end{bmatrix}$, we have, in analogy with Section 5.1.2,

$$\bar{\Psi}_j\left(\bar{\Phi}^{ETEL}_{l_\tau+h,jk}-\bar{\Phi}^{EL}_{l_\tau+h,jk}\right)\bar{\Psi}_k=\mathbf{0}$$

$$\bar{\Psi}_j\left(\bar{\Phi}^{ETEL}_{l_\kappa+h,jk}-\bar{\Phi}^{EL}_{l_\kappa+h,jk}\right)\bar{\Psi}_k=\mathbf{0}$$

$$\bar{\Psi}_j\left(\bar{\Phi}^{ETEL}_{l_\lambda+h,jk}-\bar{\Phi}^{EL}_{l_\lambda+h,jk}\right)\bar{\Psi}_k=\left[\begin{array}{cccc} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{array}\right]\left[\begin{array}{cccc} 0 & 0 & 0 & \bar{G}_{h\cdot} \\ 0 & -2\overline{g_h gg'} & \overline{g_h gg'} & 0 \\ 0 & \overline{g_h gg'} & 0 & 0 \\ \bar{G}'_{h\cdot} & 0 & 0 & 0 \end{array}\right]\left[\begin{array}{c} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{array}\right]$$

$$=\mathbf{0}$$

$$\bar{\Psi}_j\left(\bar{\Phi}^{ETEL}_{l_\theta+h,jk}-\bar{\Phi}^{EL}_{l_\theta+h,jk}\right)\bar{\Psi}_k=\left[\begin{array}{cccc} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{array}\right]\left[\begin{array}{cccc} 0 & 0 & \bar{G}'_{\cdot h} & 0 \\ 0 & -\left(\overline{\frac{\partial(gg')}{\partial\theta_h}}\right) & \left(\overline{\frac{\partial(gg')}{\partial\theta_h}}\right) & 0 \\ \bar{G}_{\cdot h} & \left(\overline{\frac{\partial(gg')}{\partial\theta_h}}\right) & -\left(\overline{\frac{\partial(gg')}{\partial\theta_h}}\right) & 0 \\ 0 & 0 & 0 & 0 \end{array}\right]\left[\begin{array}{c} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{array}\right]$$

$$=\mathbf{0}$$

where $\overline{g_h gg'}\equiv\left[n^{-1/2}\sum_i\left(\dot{g}_h\dot{g}\dot{g}'-E\left[\dot{g}_h\dot{g}\dot{g}'\right]\right)\right]_{\beta=\beta^*}$ and $\overline{\frac{\partial(gg')}{\partial\theta_h}}=\left[n^{-1/2}\sum_i\left(\overline{\frac{\partial(gg')}{\partial\theta_h}}-E\left[\frac{\partial(gg')}{\partial\theta_h}\right]\right)\right]_{\beta=\beta^*}$.

It follows that $\bar{\Psi}_j\left(\bar{\Psi}^{ETEL}_{l,jk}-\bar{\Psi}^{EL}_{l,jk}\right)\bar{\Psi}_k=\Phi^{-1}_{lh}\bar{\Psi}_j\left(\bar{\Phi}^{ETEL}_{h,jk}-\bar{\Phi}^{EL}_{h,jk}\right)\bar{\Psi}_k=0.$

6.4 Calculation of $\left(\Psi^{ETEL}_{l,jkh}-\Psi^{EL}_{l,jkh}\right)\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_h$

6.4.1 Simplification

Since $\bar{R}_{l_\theta+l}$ contributes to the n^{-2} variance only through its correlation with $\bar{\Psi}_{l_\theta+m}$, we can omit terms of $\bar{R}_{l_\theta+l}$ that are uncorrelated with $\bar{\Psi}_{l_\theta+m}$. The $O\left(n^{-2}\right)$ correlation of $\left(\Psi^{ETEL}_{l_\theta+l,jkq}-\Psi^{EL}_{l_\theta+l,jkq}\right)\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q$ with $\bar{\Psi}_{l_\theta+m}$ can be written as

$$\left(\Psi^{ETEL}_{l_\theta+l,jkq}-\Psi^{EL}_{l_\theta+l,jkq}\right)E\left[\bar{\Psi}_j\bar{\Psi}_k\right]E\left[\bar{\Psi}_q\bar{\Psi}_{l_\theta+m}\right]+\left(\Psi^{ETEL}_{l_\theta+l,jkq}-\Psi^{EL}_{l_\theta+l,jkq}\right)E\left[\bar{\Psi}_j\bar{\Psi}_q\right]E\left[\bar{\Psi}_k\bar{\Psi}_{l_\theta+m}\right]+$$
$$+\left(\Psi^{ETEL}_{l_\theta+l,jkq}-\Psi^{EL}_{l_\theta+l,jkq}\right)E\left[\bar{\Psi}_j\bar{\Psi}_{l_\theta+m}\right]E\left[\bar{\Psi}_k\bar{\Psi}_q\right].$$

Since

$$E\left[\bar{\Psi}_\cdot\bar{\Psi}'_\cdot\right]=\left[\begin{array}{cccc} 0 & 0 & 0 & 0 \\ 0 & P & P & 0 \\ 0 & P & P & 0 \\ 0 & 0 & 0 & \Sigma \end{array}\right],$$

a term of the form $E\left[\bar{\Psi}_q\bar{\Psi}_{l_\theta+m}\right]$ will be nonzero only if $q>l_\theta$. It follows that, in the sum $\left(\Psi^{ETEL}_{l,jkq}-\Psi^{EL}_{l,jkq}\right)\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q$, we only need to keep terms such that either $j>l_\theta$ or $k>l_\theta$ or $q>l_\theta$.

Letting $\Psi^{ETEL-EL}_{l_\theta+l,jkq}=\left(\Psi^{ETEL}_{l_\theta+l,jkq}-\Psi^{EL}_{l_\theta+l,jkq}\right)$, and making use of the fact that $\Psi_{l,jkq}=\Psi_{l,kjq}=\Psi_{l,qjk}=\ldots$, we have,

$$\begin{aligned}
&\sum_{j,k,q} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q 1\left(j>l_\theta \text{ or } k>l_\theta \text{ or } q>l_\theta\right)\\
=\ &\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \sum_j\sum_k\sum_{q\le l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q 1\left(j>l_\theta \text{ or } k>l_\theta\right)\\
=\ &\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \sum_j\sum_{k\le l_\theta}\sum_{q} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q 1\left(j>l_\theta \text{ or } q>l_\theta\right)\\
=\ &\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \sum_j\sum_{k\le l_\theta}\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q +\\
&+\sum_j\sum_{k\le l_\theta}\sum_{q\le l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q 1\left(j>l_\theta\right)\\
=\ &\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \sum_j\sum_{k\le l_\theta}\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \sum_{j>l_\theta}\sum_{k\le l_\theta}\sum_{q\le l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q\\
=\ &\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \sum_j\sum_{k\le l_\theta}\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \sum_{j\le l_\theta}\sum_{k\le l_\theta}\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q\\
=\ &\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \frac{1}{2}\sum_{j\le l_\theta}\sum_{k}\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q + \frac{1}{2}\sum_{j}\sum_{k\le l_\theta}\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q +\\
&+\sum_{j\le l_\theta}\sum_{k\le l_\theta}\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q\\
=\ &\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q\xi_{jk}
\end{aligned}$$

where

$$\xi_{jk}=\begin{cases} 1 & \text{if } j>l_\theta \text{ and } k>l_\theta\\ 3/2 & \text{if } \left(j>l_\theta \text{ and } k\le l_\theta\right) \text{ or } \left(j\le l_\theta \text{ and } k>l_\theta\right)\\ 3 & \text{if } j\le l_\theta \text{ and } k\le l_\theta\end{cases}$$

$\sum_j\sum_k\sum_{q>l_\theta} \Psi^{ETEL-EL}_{l_\theta+l,jkq}\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_q\xi_{jk} = \Phi^{-1}_{l_\theta+l,h}\sum_{q>l_\theta}\left(\sum_j\sum_k\left(\Phi^{ETEL}_{h,jkq}-\Phi^{EL}_{h,jkq}\right)\xi_{jk}\bar{\Psi}_j\bar{\Psi}_k\right)\bar{\Psi}_q$

6.4.2 Calculation of $\left(\Phi^{ETEL}_{h,\ j\ k\ l_\theta+q}-\Phi^{EL}_{h,\ j\ k\ l_\theta+q}\right)\xi_{jk}\bar{\Psi}_j\bar{\Psi}_k$

Noting that $\left(\dot{\phi}^{ETEL}-\dot{\phi}^{EL}\right)=\begin{bmatrix} 0\\ 0\\ \left(\tau-1\right)\dot{g}+\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{g}\dot{g}'\kappa\\ \left(\dot{\tau}-\dot{\varepsilon}\right)\dot{G}'\kappa+\dot{\tau}\dot{G}'\lambda\dot{g}'\kappa-\left(\dot{\tau}-\tau\right)\dot{G}'\lambda\end{bmatrix}$, we obtain:

(When writing the intermediate steps of the calculations, we omit the terms that will be multiplied by λ or κ after all the derivatives have been evaluated, since these terms will vanish when the true values $\lambda=0$ and $\kappa=0$ are substituted in.)

$\left(\Phi^{ETEL}_{l_\tau+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\tau+h,jk(l_\theta+q)}\right)=\mathbf{0}$

$\left(\Phi^{ETEL}_{l_\kappa+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\kappa+h,jk(l_\theta+q)}\right)=\mathbf{0}$

$\left(\Phi^{ETEL}_{l_\lambda+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\lambda+h,jk(l_\theta+q)}\right)=E^*\left[\frac{\partial^3}{\partial\beta\partial\beta'\partial\theta_q}\left(\left(\tau-1\right)\dot{g}_h\right)+\frac{\partial^3}{\partial\beta\partial\beta'\partial\theta_q}\left(\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{g}_h\dot{g}'\kappa\right)\right]=$

$$=\begin{bmatrix} 0 & 0 & 0 & E^*\left[\frac{\partial^2\dot{g}_h}{\partial\theta'\partial\theta_q}\right]\\ 0 & 0 & 0 & 0\\ 0 & 0 & 0 & 0\\ E^*\left[\frac{\partial^2\dot{g}'_h}{\partial\theta\partial\theta_q}\right] & 0 & 0 & 0\end{bmatrix}+$$

$$+\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & -2E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\varepsilon}^{2}\dot{g}_h\dot{g}\dot{g}'\right)\right] & E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\tau}\dot{g}_h\dot{g}\dot{g}'\right)\right] & 0 \\ 0 & E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\tau}\dot{g}_h\dot{g}\dot{g}'\right)\right] & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$=\begin{bmatrix} 0 & 0 & 0 & E^{*}\left[\frac{\partial^{2}\dot{g}_h}{\partial\theta'\partial\theta_q}\right] \\ 0 & -2E^{*}\left[\frac{\partial\left(\dot{g}_h\dot{g}\dot{g}'\right)}{\partial\theta_q}\right] & E^{*}\left[\frac{\partial\left(\dot{g}_h\dot{g}\dot{g}'\right)}{\partial\theta_q}\right] & 0 \\ 0 & E^{*}\left[\frac{\partial\left(\dot{g}_h\dot{g}\dot{g}'\right)}{\partial\theta_q}\right] & 0 & 0 \\ E^{*}\left[\frac{\partial^{2}\dot{g}'_h}{\partial\theta\partial\theta_q}\right] & 0 & 0 & 0 \end{bmatrix}$$

$$\left(\Phi^{ETEL}_{l_\lambda+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\lambda+h,jk(l_\theta+q)}\right)=$$

$$=E^{*}\left[\frac{\partial^{3}}{\partial\beta\partial\beta'\partial\theta_q}\left(\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{G}'_{\cdot h}\kappa\right)+\frac{\partial^{3}}{\partial\beta\partial\beta'\partial\theta_q}\left(\dot{\tau}\dot{G}'_{\cdot h}\lambda\dot{g}'\kappa\right)+\frac{\partial^{3}}{\partial\beta\partial\beta'\partial\theta_q}\left(\left(\tau-\dot{\tau}\right)\dot{G}'_{\cdot h}\lambda\right)\right]=$$

$$=\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & -E^{*}\left[\left(\frac{\partial}{\partial\theta_q}\left(\dot{\varepsilon}^{2}\dot{g}\dot{G}'_{\cdot h}+\dot{\varepsilon}^{2}\dot{G}_{\cdot h}\dot{g}'\right)\right)\right] & E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\tau}\dot{G}_{\cdot h}\dot{g}'\right)\right] & 0 \\ 0 & E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\tau}\dot{g}\dot{G}'_{\cdot h}\right)\right] & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}+$$

$$+\begin{bmatrix} 0 & 0 & 0 & 0 \\ 0 & 0 & E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\tau}\dot{g}\dot{G}'_{\cdot h}\right)\right] & 0 \\ 0 & E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\tau}\dot{G}_{\cdot h}\dot{g}'\right)\right] & 0 & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}+$$

$$+\begin{bmatrix} 0 & 0 & E^{*}\left[\frac{\partial}{\partial\theta_h}\dot{G}'_{\cdot h}\right] & 0 \\ 0 & 0 & 0 & 0 \\ E^{*}\left[\frac{\partial}{\partial\theta_h}\dot{G}_{\cdot h}\right] & 0 & -E^{*}\left[\frac{\partial}{\partial\theta_q}\left(\dot{\tau}\dot{g}\dot{G}'_{\cdot h}+\dot{\tau}\dot{G}_{\cdot h}\dot{g}'\right)\right] & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

$$=\begin{bmatrix} 0 & 0 & E^{*}\left[\frac{\partial^{2}\dot{g}'}{\partial\theta_q\partial\theta_h}\right] & 0 \\ 0 & -E^{*}\left[\frac{\partial^{2}\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & E^{*}\left[\frac{\partial^{2}\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & 0 \\ E^{*}\left[\frac{\partial^{2}\dot{g}}{\partial\theta_q\partial\theta_h}\right] & E^{*}\left[\frac{\partial^{2}\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & -E^{*}\left[\frac{\partial^{2}\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & 0 \\ 0 & 0 & 0 & 0 \end{bmatrix}$$

It follows that

$$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\tau+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\tau+h,jk(l_\theta+q)}\right)\bar{\Psi}_k\xi_{jk}=\mathbf{0}$$

$$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\kappa+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\kappa+h,jk(l_\theta+q)}\right)\bar{\Psi}_k\xi_{jk}=\mathbf{0}$$

$$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\lambda+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\lambda+h,jk(l_\theta+q)}\right)\bar{\Psi}_k\xi_{jk}=\bar{\Psi}_j\left(\frac{\partial^{3}}{\partial\theta_q\partial\beta_j\partial\beta_k}\left(\left(\tau-1\right)\dot{g}_h+\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{g}\dot{g}'\kappa\right)\right)\xi_{jk}\bar{\Psi}_k=$$

$$=\begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix}'\begin{bmatrix} 0 & 0 & 0 & \left(\frac{3}{2}\right)E^{*}\left[\frac{\partial^{2}\dot{g}_h}{\partial\theta'\partial\theta_q}\right] \\ 0 & -2\left(3\right)E^{*}\left[\frac{\partial\left(\dot{g}_h\dot{g}\dot{g}'\right)}{\partial\theta_q}\right] & \left(3\right)E^{*}\left[\frac{\partial\left(\dot{g}_h\dot{g}\dot{g}'\right)}{\partial\theta_q}\right] & 0 \\ 0 & \left(3\right)E^{*}\left[\frac{\partial\left(\dot{g}_h\dot{g}\dot{g}'\right)}{\partial\theta_q}\right] & 0 & 0 \\ \left(\frac{3}{2}\right)E^{*}\left[\frac{\partial^{2}\dot{g}'_h}{\partial\theta\partial\theta_q}\right] & 0 & 0 & 0 \end{bmatrix}\begin{bmatrix} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{bmatrix}$$

$= 0$, where the parenthesized terms (3) and $\left(\frac{3}{2}\right)$ arise from ξ_{jk}

$$\bar{\Psi}_j\left(\Phi^{ETEL}_{l_\lambda+h,jk(l_\theta+q)}-\Phi^{EL}_{l_\lambda+h,jk(l_\theta+q)}\right)\bar{\Psi}_k\xi_{jkq}=\bar{\Psi}_j\left(\frac{\partial^{3}}{\partial\theta_q\partial\beta\partial\beta'}\left(\left(\dot{\tau}-\dot{\varepsilon}\right)\dot{G}'_{\cdot h}\kappa+\dot{\tau}\dot{G}'_{\cdot h}\lambda\dot{g}'\kappa+\left(\tau-\dot{\tau}\right)\dot{G}'_{\cdot h}\lambda\right)\right)\xi_{jk}\bar{\Psi}_k=$$

$$= \left[\begin{array}{cccc} 0 & -\bar{g}'P & -\bar{g}'P & -\bar{g}'H' \end{array}\right] \left[\begin{array}{cccc} 0 & 0 & (3)\,E^*\left[\frac{\partial^2 \dot{g}'}{\partial\theta_q\partial\theta_h}\right] & 0 \\ 0 & -(3)\,E^*\left[\frac{\partial^2\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & (3)\,E^*\left[\frac{\partial^2\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & 0 \\ (3)\,E^*\left[\frac{\partial^2 \dot{g}}{\partial\theta_q\partial\theta_h}\right] & (3)\,E^*\left[\frac{\partial^2\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & -(3)\,E^*\left[\frac{\partial^2\left(\dot{g}\dot{g}'\right)}{\partial\theta_q\partial\theta_h}\right] & 0 \\ 0 & 0 & 0 & 0 \end{array}\right] \left[\begin{array}{c} 0 \\ -P\bar{g} \\ -P\bar{g} \\ -H\bar{g} \end{array}\right]$$

$$= 0$$

This implies that $\Phi^{-1}_{l_\theta+l,h}\left(\Phi^{ETEL}_{h,jkq} - \Phi^{EL}_{h,jkq}\right)\bar{\Psi}_j\bar{\Psi}_k\xi_{jk} = 0$ and thus that

$$E\left[\left(\Psi^{ETEL}_{l_\theta+l,jkh} - \Psi^{EL}_{l_\theta+l,jkh}\right)\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_h\bar{\Psi}_{l_\theta+m}\right] = 0 + o\left(n^{-2}\right).$$

7 Conclusion

$\bar{\Psi}^{ETEL}_{\cdot} = \bar{\Psi}^{EL}_{\cdot}$

$\bar{Q}^{ETEL}_{\cdot} = \bar{Q}^{EL}_{\cdot}$

$\bar{R}^{ETEL}_{\cdot} \neq \bar{R}^{EL}_{\cdot}$ but $E\left[\left(\bar{R}^{ETEL}_{l_\theta+l} - \bar{R}^{EL}_{l_\theta+l}\right)\bar{\Psi}_{l_\theta+m}\right] = o\left(n^{-2}\right)$.

In particular, the four terms entering $\left(\bar{R}^{ETEL}_{l_\theta+l} - \bar{R}^{EL}_{l_\theta+l}\right)$ have the following properties

$\left(\bar{\Psi}^{ETEL}_{l_\theta+l,j} - \bar{\Psi}^{EL}_{l_\theta+l,j}\right)\bar{Q}_j = \frac{1}{2}H_{l\cdot}\bar{g}\bar{g}'P\bar{g}$ (see Section 6.1).

$\left(\Psi^{ETEL}_{l_\theta+l,jk} - \Psi^{EL}_{l_\theta+l,jk}\right)\bar{\Psi}_k\bar{Q}_j = -\frac{1}{2}H_{l\cdot}\bar{g}\bar{g}'P\bar{g} - \Xi_{7,l}$ where $\Xi_{7,l}$ is such that $E\left[\Xi_{7,l}\bar{\Psi}_{l_\theta+m}\right] = o\left(n^{-2}\right)$ (see Section 6.2).

$\left(\bar{\Psi}^{ETEL}_{l,jk} - \bar{\Psi}^{EL}_{l,jk}\right)\bar{\Psi}_j\bar{\Psi}_k = 0$ (see Section 6.3).

$\left(\Psi^{ETEL}_{l_\theta+l,jkh} - \Psi^{EL}_{l_\theta+l,jkh}\right)\bar{\Psi}_j\bar{\Psi}_k\bar{\Psi}_h = \Xi_{8,l}$ such that $E\left[\Xi_{8,l}\bar{\Psi}^{EL}_{l_\theta+m}\right] = o\left(n^{-2}\right)$ (see Section 6.4).